A RELATION BETWEEN HOCHSCHILDT HOMOLOGY AND COHOMOLOGY FOR GORENSTEIN RINGS

MICHEL VAN DEN BERGH

Abstract. Let "HH" stand for Hochschildt (co)homology. In this note we show that for many rings A there exists $d \in \mathbb{N}$ such that for an arbitrary A-bimodule N we have $HH^i(N) = HH_{d-i}(N)$. Such a result may be viewed as an analog of Poincare duality.

Combining this equality with a computation of Soergel allows one to compute the Hochschildt homology of a regular minimal primitive quotient of an enveloping algebra of a semisimple Lie algebra, answering a question of Polo.

In the sequel the base field will be denoted by k. Let $\mathfrak{g}$ be a semisimple Lie algebra and let A be a regular minimal primitive quotient of $U(\mathfrak{g})$. The Hochschildt cohomology of A was computed by Soergel in [6] and shown to be equal to the cohomology of the corresponding flag variety. Soergel's computation is rather ingenious and makes use of the Bernstein-Beilinson theorem together with the Riemann-Hilbert correspondence. The case of singular A is still open.

After Soergel's result Patrick Polo asked whether perhaps the Hochschildt *homology* of A also coincided with the *homology* of the underlying flag manifold. It is indeed rather likely that Soergel's techniques can be adapted to this end.

In this note we answer Polo's question with a different method. Indeed we show, using elementary homological algebra, that for many algebras A (in particular those considered above) we actually have

$$HH^i(N) = HH_{d-i}(N) \tag{1}$$

for some $d \in \mathbb{N}$ and for an arbitrary A-bimodule N (see Corollary 6). This can be considered a kind of Poincare duality. In more general situations (1) may remain true provided we twist N on the right hand side by an automorphism or an invertible bimodule (see Theorem 1).

(1) maybe used to complete the results in [7] where the Hochschildt homology of a generic three dimensional Type A regular algebra was computed (see [1] for terminology). Since we show below that (1) holds for such a algebras, we now also obtain the Hochschildt cohomology without any extra work.

Our basic result is the following theorem which is easily proved.

Theorem 1. *Assume that A is an algebra such that there exists a $d \in \mathbb{N}$ with the property that $\operatorname{Ext}^i_{A^e}(A, A^e) = 0$, unless $i = d$. Put $U = \operatorname{Ext}_A(A, A^e)$ and assume in addition that U is an invertible A-bimodule. Then for every A-bimodule N we have*

$$HH^i(N) = HH_{d-i}(U \otimes_A N) \tag{2}$$

1991 *Mathematics Subject Classification.* Primary 16E40;
Key words and phrases. Hochschildt homology, Gorenstein rings.
The author is a senior researcher at the NFWO.

Proof. To prove this it is convenient to make use of the derived category. We have

$$
\begin{aligned}
&HH^i(N) = H^i(\mathrm{RHom}_{A^e}({}_{A^e}A, {}_{A^e}N)) = H^i(\mathrm{RHom}_{A^e}({}_{A^e}A, A^e) \overset{L}{\otimes}_{A^e} N) \\
&= H^i(U[-d]\overset{L}{\otimes}_{A^e} N) = H^{i-d}(U\overset{L}{\otimes}_{A^e} N) = H^{i-d}(A\overset{L}{\otimes}_{A^e}(U\overset{L}{\otimes}_A A)) = HH_{d-i}(U\otimes_A N)
\end{aligned}
$$

□

We now give several situations in which the theorem can be applied. First from [8] we recall some results concerning Koszul algebras.

Proposition 2. *Let A be a (graded) Koszul algebra of global dimension d such that $A^!$ is Frobenius. Then the hypotheses of Theorem 1 are satisfied, and furthermore we have the following precise description of U.*

Let $\phi^!$ be the graded automorphism of $A^!$ such that $(A^!)^* = A^!_{\phi^!}$. Furthermore let ϕ be the automorphism of A which is adjoint to $\phi^!$. Finally let ϵ be the automorphism of A which is multiplication by $(-)^m$ on A_m. Then

$$U = A_{\phi^{-1}\epsilon^{d+1}}$$

In particular (2) *specializes to*

$$HH^i(N) = HH_{d-i}({}_{\phi\epsilon^{d+1}}N)$$

The simplest non-trivial Koszul algebras of finite global dimension are those of global dimension three. These have been studied extensively in [1, 2, 3]. Recall that they have a presentation $k\langle x_1, x_2, x_3\rangle/(f_1, f_2, f_3)$ in which $x = (x_1, x_2, x_3)^t$, $f = (f_1, f_2, f_3)^t$ can be chosen in such a way that $f = Mx$, $x^t M = (Qf)^t$ for 3×3-matrices M and Q, respectively with linear an scalar entries. It is shown in [8] that in this case the automorphism ϕ is given in terms of the generators by $x \mapsto Q^t x$. In particular if A is a type A algebra (see [1]) then (1) holds with $d = 3$.

Let us now exhibit another situation in which Theorem 1 applies and which occurs frequently in practice.

Proposition 3. *Let A be a k-algebra equipped with a k-linear ascending filtration $(F_n A)_{n\geq 0}$ such that $\mathrm{gr}_F A$ is commutative of finite type over k. Assume that $\mathrm{gr}\, A$ is Gorenstein of dimension d. Then the hypotheses of Theorem 1 hold and furthermore U carries an A-bimodule filtration such that $\mathrm{gr}\, U$ is equal to $\omega^{-1}_{\mathrm{gr}\, A/k}$, the inverse of the dualizing module of $\mathrm{gr}\, A$.*

Proof. We use the standard spectral sequence

$$\mathrm{Ext}^i_{(\mathrm{gr}\, A)^e}(\mathrm{gr}\, A, (\mathrm{gr}\, A)^e) \Rightarrow \mathrm{Ext}^i_{A^e}(A, A^e) \tag{3}$$

which converges because we have positive filtrations.

Let C is a commutative Cohen-Macaulay ring of dimension d. From the adjunction formula one has

$$\mathrm{RHom}_{C^e}(C, \omega_{C^e}) = \omega_C[-d]$$

Assume now that C is Gorenstein. Then $\omega_{C^e} = \omega_C \otimes_k \omega_C$ and ω_C are invertible and hence

$$\mathrm{RHom}_{C^e}(C, \omega_{C^e}) = \mathrm{RHom}_{C^e}(C, C^e) \otimes_{C^e} \omega_{C^e}$$

Combining this yields

$$\mathrm{RHom}_{C^e}(C, C^e) = \omega_C[-d] \otimes_{C^e} \omega^{-1}_{C^e} = \omega^{-1}_C[-d]$$

Applying this with $C = \operatorname{gr} A$ shows that (3) degenerates and we obtain

$$\operatorname{RHom}_{A^e}(A, A^e) = U[-d]$$

where U lives in one degree and where, from a functoriality argument using (3), one obtains that the filtration on U is an A^e filtration, hence an A bimodule filtration. Furthermore for this filtration, $\operatorname{gr} U = \omega_C^{-1}$. It now suffices to apply lemma 4 below to deduce that U is invertible. □

Lemma 4. *Assume that A is a ring equipped with a positive ascending filtration and let U be an A-bimodule, equipped with a left limited A-bimodule filtration. If $\operatorname{gr} U$ is an invertible $\operatorname{gr} A$-bimodule then U is an invertible A-bimodule.*

Proof. We have to show [4, §1] that U is left and right projective and that the canonical maps

$$A \to \operatorname{End}_A({}_A U) \tag{4}$$

$$A \to \operatorname{End}_A(A_U) \tag{5}$$

are surjective. The fact that U is projective is standard (see [5, Cor. D.VII.6]). Therefore let us concentrate on (4). (5) is similar. We use the standard spectral sequence

$$\operatorname{Ext}^i(\operatorname{gr}({}_A U), \operatorname{gr}({}_A U)) \Rightarrow \operatorname{Ext}^i_A(U, U)$$

which is compatible with (4) and which obviously degenerates. From this we deduce that

$$\operatorname{gr} \operatorname{End}_A({}_A U) = \operatorname{End}_A(\operatorname{gr}({}_A U))$$

Combining this with the hypotheses that $\operatorname{gr} U$ is invertible yields that

$$\operatorname{gr} A \to \operatorname{gr} \operatorname{End}_A({}_A U)$$

is surjective. Hence (4) is also surjective. □

We use the following lemma below.

Lemma 5. *Let the notations as in the previous lemma. Assume that $\operatorname{gr} U$ is left free of rank one and let θ be the automorphism of $\operatorname{gr} A$ such that $\operatorname{gr} U = (\operatorname{gr} A)_\theta$. Then there exists a filtered automorphism ψ of A such that $U = A_\psi$ and $\operatorname{gr} \psi = \theta$.*

Proof. Let $\bar{u}$ be a generator of $\operatorname{gr} U$ such that

$$\bar{u} y = \theta(y) \bar{u} \tag{6}$$

and let u be a lifting to U. It is easy to see that u is a basis for U. Thus there is an automorphism ψ of A such that $ux = \psi(x)u$. If we take the image of this equality in $\operatorname{gr} A$ we find $\bar{u}\bar{x} = \overline{\psi(x)}\bar{u}$. Comparing with (6) yields $\theta(\bar{x}) = \overline{\psi(x)}$. From this we easily deduce that ψ is filtered, and $\operatorname{gr} \psi = \theta$. □

Corollary 6. *Assume that A is equipped with a filtration $(F_n A)_{n \in \mathbb{N}}$ such that $F_0 A = k$ and such that $\operatorname{gr} A$ is commutative and generated by $\operatorname{gr}_1 A$. Put $\mathfrak{g} = \operatorname{gr}_1 A$ with Lie algebra structure deduced from the commutator on A. Assume furthermore that $\operatorname{gr} A$ is Gorenstein and $[\mathfrak{g}, \mathfrak{g}] = \mathfrak{g}$. Then*

$$HH^i(N) = HH_{d-i}(N)$$

Proof. Since $\operatorname{gr} A$ is graded local we have $\omega_A = A$, as A-bimodules and hence we know from lemma 5 that $H^i(N) = H_{d-i}({}_{\psi^{-1}}N)$ for some filtered automorphism of A such that $\operatorname{gr}\psi = \mathrm{id}_{\operatorname{gr} A}$. Let us consider $\mathfrak{g}$ as a subspace of $F_1 A$. For $x, y \in \mathfrak{g}$ we have

$$yx - xy = [y, x] + \alpha \tag{7}$$

for some $\alpha \in k$. We also have $\psi(x) = x + \beta(x)$, where $\beta \in \mathfrak{g}^*$. Applying ψ to (7) yields

$$yx - xy = [y, x] + \beta([y, x]) + \alpha$$

and thus $\beta \mid [\mathfrak{g}, \mathfrak{g}] = 0$. Hence $\beta = 0$ and we are done. □

References

[1] M. Artin and W. Schelter, *Graded algebras of global dimension 3*, Adv. in Math. **66** (1987), 171–216.

[2] M. Artin, J. Tate, and M. Van den Bergh, *Some algebras associated to automorphisms of elliptic curves*, The Grothendieck Festschrift, vol. 1, Birkhäuser, 1990, pp. 33–85.

[3] ———, *Modules over regular algebras of dimension 3*, Invent. Math. **106** (1991), 335–388.

[4] A. Fröhlich, *The Picard group of non-commutative rings*, Trans. Amer. Math. Soc. **180** (1973), 1–45.

[5] C. Nastacescu and F. Van Oystaeyen, *Graded ring theory*, North-Holland, 1982.

[6] W. Soergel, *The Hochschild cohomology of regular maximal primitive quotients of enveloping algebras of semisimple Lie algebras*, preprint Universität Freiburg, 1995.

[7] M. Van den Bergh, *Non-commutative homology of some three dimensional quantum spaces*, J. K-theory (1994), 213–230.

[8] ———, *Existence theorems for dualizing complexes over non-commutative graded and filtered rings*, preprint Limburgs Universitair Centrum, 1996.

Departement WNI, Limburgs Universitair Centrum, Universitaire Campus, Building D, 3590 Diepenbeek, Belgium

E-mail address, M. Van den Bergh: vdbergh@luc.ac.be